201-14216



The Officer KOlena MillFe to.comP at 171 M2002 08:26,35 PM

To:

Rtk Chem/DC/USEPA/US@EPA, oppt.ncic@epamail.epa.gov

cc:

Subject: HPV Filing

Attached is the HPV Filing for Isodecyl Diphenyl Phosphate, CAS 29761-21-5.

Alan J. Olson Director of Technology Ferro Corp.

IDP HVP cl 123002.doc IDPHPVtestplan 123002.doc

2003 JAN - 2 PH 2: 54

OPPT NOIC

December 30, 2002 Via email and FedEx

Mr. Oscar Hernandez
Director, Risk Assessment Division (7403M)
U. S. Environmental Protection Agency
1200 Pennsylvania Ave. NW
Washington, DC 20460

Attn: Chemical Right-To-Known Program Re: HPV Challange Isodecyl Diphenyl Phosphate CAS Registry Number 29761-21-5

Dear Mr. Hernandez:

As part of Ferro Corporation's commitment under EPA's High Production Volume (HPV) Challenge Program, Ferro is pleased to submit its proposed testing approach for Isodecyl Diphenyl Phosphate. This submission consists of this cover letter, a Test Plan, and robust summaries for the existing studies available on Isodecyl Diphenyl Phosphate.

Ferro understands that this Test Plan will be posted on the Internet and subject to a 120-day comment period. Ferro further understands that all comments by EPA or received by EPA will be forwarded to Ferro for consideration. In the event that additional information on existing studies of Isodecyl Diphenyl Phosphate

becomes available to Ferro during the 120-day comment period, Ferro may wish to amend its submission.

This submission is also being sent electronically to the following e-mail addresses:

oppt.ncic@epa.gov chem.rtk@epa.gov.

Thank you for your cooperation in this matter. If EPA requires any additional information about this submission, please contact me at 216-750-6696 or olsona@ferro.com.

Sincerely,

Alan J. Olson, P.E. Director of Technology

U.S. EPA HIGH PRODUCTION VOLUME CHEMICAL VOLUNTARY TESTING PROGRAM

TEST PLAN

ISODECYL DIPHENYL PHOSPHATE

Submitted by:

FERRO CORPORATION CLEVELAND, OHIO

December 2002

INTRODUCTION

Isodecyl diphenyl phosphate, CAS Registry Number 29761-21-5, is a flame retardant for most commercial resins including polyvinyl chloride and its copolymers, polyvinyl acetate and acrylics. Isodecyl diphenyl phosphate is a clear, odorless liquid with the following physical properties:

Vapor pressure 0.09 kPa @ 28° C Volatility 1.6% w/w EPA method 24 Viscosity 21.9 mPa @ 25°C Solubility 0.75 mg/L @ 25°C.

TEST PLAN RATIONALE

At this time, information available on the environmental effects, ecotoxicity and some health effects of isodecyl diphenyl phosphate cannot be documented or is judged to be not reliable according to the standards specified by Klimisch (Regulatory Toxicology and Pharmacology, 25, 1-5, 1997) or the EPA High Production Volume Challenge Program Guidelines for Determining the Adequacy of Existing Data (http://www.epa.gov/chemrtk/datadfin.htm). The exceptions to this are a well-conducted and reported 90-day repeated dose oral toxicity study in rats and an in vitro mutation study conducted in mouse lymphoma cells. Summaries of those studies are included as Appendix 1 to this submission. Accordingly, Ferro Corporation commits to generating data, listed in Table 1, necessary to meet address HPV Endpoints. Since an adequate 90day repeat-dose oral toxicity study in rats has been completed for isodecyl diphenyl phosphate which established an effect level for the test compound and identified target organs for toxicity, no further repeat-dose testing is necessary. Furthermore, since the 90-day study included gross and microscopic examinations of reproductive tissues from male and female animals, and no compound-related adverse effect was produced in any reproductive organ including the testes, developmental toxicity testing is proposed to meet the HPV requirements for reproductive/developmental toxicity endpoints.

Because mammalian cell (L5178Y mouse lymphoma cells) testing for genetic toxicity has shown isodecyl diphenyl phosphate to be cytotoxic but not genotoxic or mutagenic, Ferro proposes no additional gene mutation toxicity testing for this compound. Given the degree of cytotoxicity observed in mammalian cells exposed to isodecyl diphenyl phosphate, bioavailability is not an issue and it is therefore unlikely that an Ames test would add any meaningful information to what exists insofar as gene mutation. Ferro proposed to conduct an *in vitro* chromosome aberration test to complete the genetic toxicity portion of the SIDS HPV screen.

Ferro Corporation is committed to providing EPA with reliable data necessary to complete the SIDS screening matrix for the HPV voluntary challenge; however, Ferro

Corporation is also committed to judicious use of research animal resources. To this end Ferro Corporation will continue to attempt to obtain adequate documentation on existing studies of isodecyl diphenyl phosphate. To the extent that this documentation becomes available to Ferro, the HPV Test Plan submitted herein may be altered to reflect reliance on existing studies.

TEST PLAN: ISODECYL DIPHENYL PHOSPHATE

Table 1 lists the HPV testing planned by Ferro Corporation for isodecyl diphenyl phosphate.

Table 1: ISODECYL DIPHENYL PHOSPHATE HPV TEST PLAN

HPV DATA	PROPOSED DATA DEVELOPMENT METHOD
ENDPOINT	
1. CHEMISTRY	
Melting Point	OECD Test Guideline 102
Boiling Point	OECD Test Guideline 103
Vapor Pressure	OECD Test Guideline 104
Water Solubility	OECD Test Guideline 105
Partition Co-	OECD Test Guideline 107
Efficient	
2. ENVIRON-	
MENTAL FATE	
Photodegradation	Estimate/model
Hydrolysis	OECD Test Guideline 111
(Stability in Water)	
Biodegradation	OECD Test Guideline 301
Fugacity	Fugacity Level III Modeling
3. HEALTH EFFECTS	
Acute Toxicity	"Up and Down Method" for Acute Oral Toxicity: OECD
	Health Effects Test Guideline 425 possibly supplemented by in
	vitro testing for dose-range finding
Repeat Dose Toxicity	Adequate repeat-dose study exists; no additional systemic
Repro-Develop.	toxicity testing planned.
Toxicity	Developmental toxicity study planned: OECD Health Effects
	Test Guideline 414
Genetic Toxicity	In vitro mammalian cell mutation study exists; no additional
	gene mutation testing planned.
	In vitro Chromosome Aberration Study planned (OECD 473)
4. ECOTOXICITY	
Fish	Acute Toxicity to Fish: OECD Test Guideline 203
Daphnia	Acute Toxicity to Aquatic Invertebrates: OECD Test Guideline 202
Algae	Acute Toxicity to Aquatic Plants (Algae): OECD Test Guideline 201

APPENDIX 1

ROBUST SUMMARY OF GENETIC TOXICITY AND REPEATED DOSE MAMMALIAN TOXICITY TESTING COMPLETED ON ISODECYL DIPHENYL PHOSPHATE

GENETIC TOXICITY

Test material: Isodecyl diphenyl phosphate (Lot QH-28641)

Type: In vitro mammalian cell mutation

Cell type: Fischer mouse lymphoma L5178Y derived

Metabolic

activation: Male F-344 rat liver 9000 x G supernatant, Arochlor 1254 induced

Assay run with and without activation

Solubility, cytotoxicity

determination: Solubility and toxicity were determined with 4-hour incubation

followed by 24-hr expression times in at least 4 dose levels bracketing

the concentrations used in definitive testing.

Number

of concentrations

evaluated: Posit

Positive control, negative control, vehicle control, 5 test concentrations

with and without activation

Results: No evidence for mutagenic activity in the presence or absence of

exogenous metabolic activation. Cytotoxicity was produced in the highest concentrations of isodecyl diphenyl phosphate tested with

activation.

Reliability: Reliable

GLP: Work conducted prior to inception of GLP regulations

Reference: Litton Bionetics Inc. report 20989, "Mutagenicity Evaluation of S-148

BO-78-85 in the Mouse Lymphoma Forwatrd Mutation Assay" Kensington, MD., August, 1978. D. Matheson, Ph.D., Author.

REPEAT DOSE MAMMALIAN TOXICITY

Oral Toxicity

Test material: Isodecyl diphenyl phosphate (Lot DC 5A83) 98.4% purity)

Type: Repeat-dose oral – dietary admixture

Species: Rat

Strain: Sprague-Dawley

Sex: Female and male

Number of animals

per dose level: 30, weight range: males 106-143 at study initiation

females: 104-133 at study imitiation

Number of dose

levels: Three plus untreated control

140 ppm in diet 1400 ppm in diet 7000 ppm in diet

Administration: Daily for 90 days

Observations: Clinical observation

Food consumption Body weight

Mortality

Clinical pathology at mid-study and termination

hematology serum chemistry urinalysis

Gross and microscopic pathology

Organ weights

Electron microscopy

Analysis of test material and dietary mixtures

Results: Survival was unaffected by treatment. Dose-related male and female

reduced body weight gain (slightly >10% in high dose groups) and reduced food consumption. Lymphocytopenia, decreased red cell indices, increased gamma-GT, as well as other indicators of liver cytotoxicity and/or function, hepatocellular hypertrophy and hyperplasia. No treatment-related changes in any organ or tissue

except liver.

Reliability:

Reliable

GLP:

Work conducted prior to inception of GLP regulations

Reference:

Monsanto Company Environmental Health Laboratory Report number 820160, "Subchronic Study of Santicizer 148 Plasticizer Administered in the Diet to Albino Rats" St. Louis, MO., February, 1986, M. W.

Naylor, Study Director